

**Yellow****Weight 12 kg****Yellow**

HR 90 to 150

SBP 94 - 102

Resp 22 to 30

ET tube 4.5 mm uncuffed

Medication	Dose per kg	Dose	Supplied as	ml to give
Adenosine (IV, avoid IO)	0.1 mg/kg	<b>1.2 mg</b>	3 mg/ml	<b>0.4 ml Rapid</b>
	Repeat 0.2 mg/kg	<b>2.4 mg</b>		<b>0.8 ml Rapid</b>
Albuterol (Nebulized)	N/A	<b>2.5 mg</b>	2.5 mg in 3 ml (0.083%) Premixed ampule	<b>1 premixed ampules</b>
Amiodarone IV/IO	5 mg/kg	<b>60 mg</b>	50 mg/ml	<b>1.2 ml</b>
	Repeat up to 15 mg/kg	<b>Up to 180 mg</b>		<b>Up to 3.6 ml</b>
Atropine IV/IO	0.02 mg/kg Repeat once	<b>0.24 mg</b>	0.1 mg/ml	<b>2.4 ml</b>
Bicarb 8.4% IV/IO	1 mEq/kg	<b>12 mEq</b>	1 mEq/ml Give slowly	<b>12 ml</b>
Calcium Cl 10% IV/IO	20 mg/kg	<b>240 mg</b>	100 mg/ml	<b>2.4 ml Slowly</b>
Charcoal, Activated PO (No sorbitol)	1 g/kg	<b>12 g</b>	25 g/120 ml	<b>60 ml</b>
D25W IV/IO	1 g/kg 4 ml/kg	<b>48 ml</b>	Dilute D50W 24 ml with SW 24 ml	<b>48 ml</b>
Diazepam IV/IO	0.2 mg/kg	<b>2.4 mg</b>	5 mg/ml	<b>0.5 ml</b>
Diazepam PR	0.5 mg/kg	<b>6 mg</b>	5 mg/ml	<b>1.2 ml</b>
Diphenhydramine PO	1 mg/kg	<b>12.5 mg</b>	12.5 mg/5 ml po	<b>5 ml</b>
Diphenhydramine IV/IO/IM	1 mg/kg	<b>12 mg</b>	50 mg/ml	<b>0.3 ml</b>
Epinephrine 1:10,000 IV/IO	0.01 mg/kg	<b>0.12 mg</b>	0.1 mg/ml (1:10,000)	<b>1.2 ml</b>
Epinephrine 1:1000 (IM/SC)	0.01 mg/kg	<b>0.12 mg</b>	1 mg/ml (1:1000)	<b>0.1 ml</b>

**Yellow****Weight 12 kg****Yellow****HR 90 to 150****SBP 94 - 102****Resp 22 to 30****ET tube 4.5 mm uncuffed**

<b>Medication</b>	<b>Dose per kg</b>	<b>Dose</b>	<b>Supplied as</b>	<b>ml to give</b>
Epinephrine (Racemic 2.25%) Nebulized	N/A	<b>0.5 ml</b>	0.5 ml Dilute in 3-5 ml NS and nebulize	<b>0.5 ml</b>
Flumazenil IV/IO	0.01 mg/kg	<b>0.12 mg</b>	0.1 mg/ml	<b>1.2 ml</b>
Fosphenytoin IV/IO/IM	20 mg PE/kg (PE= Phenytoin Equivalents)	<b>240 mg</b>	50 mg PE/ml Dilute with NS and give IV over 7 min Do NOT dilute for IM	<b>4.8 ml</b> (predilutional)
Furosemide IV/IO/IM	1 mg/kg	<b>12 mg</b>	10 mg/ml	<b>1.2 ml</b>
Lidocaine 2% IV/IO	1 mg/kg	<b>12 mg</b>	20 mg/ml	<b>0.6 ml</b>
Lorazepam IV/IO/IM	0.1 mg/kg	<b>1.2 mg</b>	2 mg/ml	<b>0.6 ml</b>
Magnesium Sulfate IV/IO	25-50 mg/kg	<b>300-600 mg</b>	500 mg/ml	<b>0.6-1.2 ml</b> Slowly
Mannitol 20% IV/IO	1 g/kg	<b>12 gm</b>	20 g/100 ml	<b>60 ml</b>
Midazolam IV/IO/IM	0.1 mg/kg	<b>1.2 mg</b>	1 mg/ml	<b>1.2 ml</b>
Naloxone IV/IO/IM/IN/ET	0.1 mg/kg	<b>1.2 mg</b>	0.4 mg/ml Or 1 mg/ml	<b>3 ml</b> Or <b>1.2 ml</b>
Phenobarbital IV/IO/IM	20 mg/kg	<b>240 mg</b>	130 mg/ml Dilute with NS Give over 20 min	<b>1.8 ml</b>
Phenytoin IV/IO	20 mg/kg	<b>240 mg</b>	50 mg/ml Dilute with NS Give over 20 min	<b>4.8 ml</b> (predilutional)
Procainamide IV/IO	15 mg/kg	<b>180 mg</b>	100 mg/ml	<b>1.8 ml</b>

**Yellow****Weight 12 kg****Yellow**

HR 90 to 150

SBP 94 - 102

Resp 22 to 30

ET tube 4.5 mm uncuffed

**Medications for Rapid Sequence Intubation**

Medication	Dose per kg	Dose	Supplied as	ml to give
<b>INDUCTION AGENTS</b>				
Etomidate	0.3 mg/kg	<b>3.6 mg</b>	2 mg/ml	<b>1.8 ml</b>
<b>PARALYZING AGENTS</b>				
Rocuronium	0.6 mg/kg	<b>7.2 mg</b>	10 mg/ml	<b>0.7 ml</b>
Succinylcholine	2 mg/kg	<b>24 mg</b>	20 mg/ml	<b>1.2 ml</b>
Vecuronium	0.1-0.2 mg/kg	<b>1.2-2.4 mg</b>	1 mg/ml 10mg/vial mixed with 10 mL NS	<b>1.2-2.4 ml</b>

**Fluid administration**

<b>Fluid bolus</b>	<b>240 ml</b>	(20 ml/kg)
<b>Blood</b>	<b>120 ml</b>	(10 ml/kg)

**Maintenance IV fluid****IV rate = 44 ml/hr****Electrical countershock****Defibrillation****24 joules** (2 j/kg)**Then 48 joules** (4 j/kg)**Synchronized****6 joules** (0.5 j/kg)**Then 12 joules** (1 j/kg)